arXiv:1303.5205v2 [math.CO] 23 Apr 2013

THE ERDŐS-HAJNAL CONJECTURE FOR PATHS AND ANTIPATHS

NICOLAS BOUSQUET, AURÉLIE LAGOUTTE, AND STÉPHAN THOMASSÉ

Abstract. We prove that for every k, there exists $c_k > 0$ such that every graph G on n vertices not inducing a path P_k and its complement $\overline{P_k}$ contains a clique or a stable set of size n^{c_k}.

An n-graph is a graph on n vertices. A class $\mathcal{C}$ of graphs (i.e. closed under induced subgraphs) is said to satisfy the *Erdős-Hajnal property* if there exists some $c > 0$ such that every n-graph of $\mathcal{C}$ contains a clique or a stable set of size n^c. The Erdős-Hajnal conjecture [6] asserts that every strict class of graphs satisfies the Erdős-Hajnal property, see [4] for a survey. This fascinating question is even open for graphs not inducing a cycle of length five. When excluding a single graph H, Alon, Pach and Solymosi showed in [2] that it suffices to consider prime H (i.e. without nontrivial modules). A natural approach is then to study classes of graphs with intermediate difficulty, hoping to get a proof scheme which could be extended. A natural prime candidate to forbid is certainly the path. Chudnovsky and Zwols studied the class $\mathcal{C}_k$ of graphs not inducing the path P_k on k vertices nor $\overline{P_k}$. They proved the Erdős-Hajnal property for P_5 and $\overline{P_6}$-free graphs [3]. This was extended for P_5 and $\overline{P_7}$-free graphs by Chudnovsky and Seymour [5]. We show in this note that for every fixed k, the class $\mathcal{C}_k$ satisfies the Erdős-Hajnal property. An n-graph is an *ε-stable set* if it has at most $\varepsilon\binom{n}{2}$ edges. The complement of an ε-stable set is an *ε-clique*. A stronger version of the following result was proved by Rödl [12]:

Theorem 1. *For every positive integer k and every $\varepsilon > 0$, there exists $\delta > 0$ such that every n-graph G satisfies one of the following:*

- *G induces all graphs on k vertices.*
- *G contains an ε-stable set of size at least δn.*
- *G contains an ε-clique of size at least δn.*

Proof. (sketch) Note that by choosing δ small enough, we can assume that G is arbitrarily large, since a single vertex is certainly an ε-stable set. Let k' and k'' be two integers such that $\max(k, 2/\varepsilon) \ll k' \ll k''$. Select now some $\varepsilon' > 0$ such that $\varepsilon' \ll \min((1/k'')^2, (\varepsilon/2)^k)$. Applying Szemerédi's lemma, there exists M for which every large enough graph has an ε'-regular partition with at least k'' parts and at most M parts. We consider such an ε'-regular partition P of G. Since ε' is small enough, there are k'' classes of P forming pairwise ε'-regular pairs. Since $k'' \gg k'$ (precisely, providing that k'' is at least $\text{Ramsey}(k', k', k')$), we can find k' of them with all pairwise densities, either less than $\varepsilon/2$, or between $\varepsilon/2$ and $1 - \varepsilon/2$, or more than $1 - \varepsilon/2$. The first case gives an ε-stable set of size δn, where δ is k'/M. The last case gives an ε-clique of size δn. Since $k \leq k'$, the intermediate case provides k parts such that all pairs are ε'-regular with densities between $\varepsilon/2$ and $1 - \varepsilon/2$. Thus one can induce all possible graphs by choosing one vertex in each of the parts since $\varepsilon' \ll (\varepsilon/2)^k$. □

Note that another proof of Theorem 1 was then provided by Fox and Sudakov [9], with no use of the regularity lemma, consequently giving a better constant $\delta = 2^{-15k(\log(1/\varepsilon))^2}$.

In a graph G, a *complete ℓ-bipartite graph* is a pair of disjoint subsets X, Y of vertices of G, both of size ℓ and inducing all edges between X and Y. We define similarly *empty ℓ-bipartite graph* when there is no edge between X and Y. Observe that we do not require any condition inside X or inside Y. Erdős, Hajnal and Pach proved in [7] that for every strict class $\mathcal{C}$, there exists some $c > 0$ such that every n-graph in $\mathcal{C}$ contains an empty or complete n^c-bipartite graph. This "half" version of

the conjecture was improved to a "three quarter" version by Fox and Sudakov [8], where they show the existence of a polynomial clique or empty bipartite graph. It was proved that getting linear complete or empty bipartite graphs is enough to prove the full version:

Theorem 2 ([1, 10])**.** *If $\mathcal{C}$ is a class of graphs for which there exists $c > 0$ such that every graph G of $\mathcal{C}$ has an empty or complete $c.n$-bipartite graph, then $\mathcal{C}$ satisfies the Erdős-Hajnal property.*

Proof. (sketch) Let $c' > 0$ such that $c^{c'} \geq 1/2$. We prove by induction that every graph G of $\mathcal{C}$ induces a P_4-free graph of size $n^{c'}$. By our hypothesis on $\mathcal{C}$, there exists, say, a complete $c.n$-bipartite graph X, Y in G. Applying the induction hypothesis inside both X and Y, we form a P_4-free graph on $2(c.n)^{c'} \geq n^{c'}$ vertices. The Erdős-Hajnal property of $\mathcal{C}$ follows from the fact that every P_4-free $n^{c'}$-graph has a clique or a stable set of size at least $n^{c'/2}$. □

Before going to our main result, we state an intermediate result. If $d(x)$ is the degree of the vertex x, its *closed degree* $\overline{d(x)}$ is $d(x)+1$, corresponding to the size of its *closed neighborhood* $N[x]$, that is to say $\{x\} \cup \{y | xy \in E\}$.

Theorem 3. *Let $n \geq 2$ be an integer, $c, \varepsilon \in [0, 1]$ and G be a connected n-graph. Assume that G contains no empty $c.n$-bipartite subgraph and that every vertex x has closed degree $\overline{d(x)} \leq \varepsilon.n$. Then for every vertex x, there exists a path P_k starting in x with $k \geq \frac{1}{2(\varepsilon+c)}$.*

Proof. We follow the lines of Gyárfás' proof of the χ-boundedness of P_k-free graphs, see [11]. First assume that $3c+\varepsilon \geq 1$. In particular, $\frac{1}{2(\varepsilon+c)} \leq \frac{3}{2}$. To conclude, observe that every vertex is incident to a P_2 (an edge) in a connected graph of size at least 2. From now on, assume that $3c + \varepsilon < 1$. Let x be a vertex, U be the set $V \setminus N[x]$, and C_1 be the largest connected component of $G[U]$. We distinguish three cases:

- if $|C_1| \geq (1 - \varepsilon - c).n$: let $y \in N[x]$ having a neighbor in C_1, and $G' = G[\{y\} \cup C_1]$. Apply the induction hypothesis to G' containing no empty $c'.n'$-bipartite subgraph with $n' = |G'| \geq (1 - \varepsilon - c).n$ and $c' = c/(1 - \varepsilon - c)$. Moreover, every vertex x of G' has closed degree $\overline{d(x)} \leq \varepsilon'.n'$ with $\varepsilon' = \varepsilon/(1 - \varepsilon - c)$. Then, by induction, there exists a path $P_{k'}$ starting in y with $k' \geq \frac{1}{2(\varepsilon'+c')} = \frac{1}{2(\varepsilon+c)} - \frac{1}{2}$. Augmenting the path by x gives a path of length $k' + 1 \geq \frac{1}{2(\varepsilon+c)}$.
- if $c.n \leq |C_1| \leq (1-\varepsilon-c).n$: then $(C_1, U \setminus C_1)$ is an empty $c.n$-bipartite graph, a contradiction.
- otherwise $|C_1| \leq c.n$: then U is divided into small connected components that we can split into two parts, each of size at least $c.n$ (this is possible because the total size of U is at least $3c.n$, due to $3c + \varepsilon < 1$). This forms an empty $c.n$-bipartite graph, a contradiction.

□

We now have all the tools for our main result:

Theorem 4. *For every k, there is a $c_k > 0$, such that every graph in $\mathcal{C}_k$ contains an empty or complete $c_k.n$-bipartite graph. Thus, by Theorem 2, the class $\mathcal{C}_k$ satisfies the Erdős-Hajnal property.*

Proof. Let $\varepsilon > 0$ be some small value. By Theorem 1, there exists $\delta > 0$ such that every graph G not inducing P_k or $\overline{P_k}$ does contain an ε-stable set or an ε-clique of size at least δn. Free to consider the complement of G, we can assume that G contains an ε-stable set S of size δn. We start by deleting in S all the vertices with degree in S more than $2\varepsilon s$ where s is the size of S. Since the average degree in S is at most εs, we do not delete more than half of the vertices. We still call S the remaining subgraph which is a 2ε-stable set of size $s \geq \delta n/2$ with maximum degree at most $2\varepsilon s$. Let $c \in [0, 1]$ be a small constant. If S contains no empty $c.s$-bipartite subgraph, we apply Theorem 3 and get a path $P_{k'}$ with $k' \geq \frac{1}{2(2\varepsilon+c)}$. Choosing ε and c small enough leads to a contradiction to G not inducing a P_k. Thus G contains an empty $c\delta/2.n$-bipartite graph. □

References

[1] N. Alon, J. Pach, R. Pinchasi, R. Radoičić and M. Sharir, Crossing patterns of semi-algebraic sets. *Journal of Combinatorial Theory Series A*, 111(2):310–326, 2005.

[2] N. Alon, J. Pach, and J. Solymosi, Ramsey-type theorems with forbidden subgraphs. *Combinatorica*, 21:155–170, 2001.

[3] M. Chudnovsky and Y. Zwols, Large cliques or stable sets in graphs with no four-edge path and no five-edge path in the complement. *Journal of Graph Theory*, 70:449–472, 2012.

[4] M. Chudnovsky, The Erdős-Hajnal Conjecture - A Survey. preprint, 2013.

[5] M. Chudnovsky and P. Seymour, Excluding paths and antipaths. preprint, 2012.

[6] P. Erdős and A. Hajnal, Ramsey-type theorems. *Discrete Applied Mathematics*, 25:37–52, 1989.

[7] P. Erdős, A. Hajnal and J. Pach, Ramsey-type theorem for bipartite graphs. *Geombinatorics*, 10:64–68, 2000.

[8] J. Fox and B. Sudakov, Density theorems for bipartite graphs and related Ramsey-type results. *Combinatorica*, 29:153–196, 2009.

[9] J. Fox and B. Sudakov, Induced Ramsey-type theorems. *Advances in Mathematics*, 219:1771–1800, 2008.

[10] J. Fox and J. Pach, Erdős-Hajnal-type results on intersection patterns of geometric objetcs *Horizon of Combinatorics* (G.O.H. Katona et al., eds.), Bolyai Society Studies in Mathematics, Springer, 79–103, 2008.

[11] A. Gyárfás, Problems from the world surrounding perfect graphs. *Zastos. Mat.*, 413–441, 1987.

[12] V. Rödl, On universality of graphs with uniformly distributed edges. *Discrete Mathematics*, 59:125–134, 1986.

Université Montpellier 2 - CNRS, LIRMM, 161 rue Ada, 34392 Montpellier, France
E-mail address: <bousquet@lirmm.fr>

LIP, UMR 5668 ENS Lyon - CNRS - UCBL - INRIA, Université de Lyon, France.
E-mail address: <aurelie.lagoutte,stephan.thomasse@ens-lyon.fr>